BLUE LIGHT THERAPY FOR ACNE - FOREVER CLEARTM

Treatment Instructions

BLUE LIGHT FOR ACNE PRE TREATMENT INSTRUCTIONS

3-6 Weeks Before

- **Pre-treatment topical medications**. For certain skin types it may be necessary to pre-treat with Retin-A and Hydroquinone for 4-6 weeks prior to scheduled treatment.
- No sun exposure, tanning beds or self-tanning products. Avoid tanning for 3 weeks prior
 to procedure and for 3 months after the IPL procedure. If tanned or sunburned procedure
 may be canceled for your safety.
- **Avoid harsh skincare products.** Avoid products or treatments that irritate the skin for two weeks prior to treatment, such as chemical peels, waxing, and chemical depilatories.
- Tell your provider if you are prone to cold sores. If you are prone to cold sores you will be
 placed on an anti-viral medication. Please take medication as instructed and complete
 entire course.

3 Days Before

• **Temporarily stop all strong topical skincare products**. Skincare products including: Retin-A, retinoids, topical antibiotics, exfoliants, hydroquinone, and benzyl peroxide should be stopped about 3 days prior to blue light therapy. Use only mild soap and sunscreen. If you have been on antibiotics oral/topical up to 7 days prior to therapy please inform our staff.

Day of Procedure

- Bring a wide brim hat to wear on the day of your procedure. Please arrive with fully cleansed skin: no makeup, moisturizers, perfume, cologne, sunscreen or any other products on the skin.
- Notify us of any medication allergies, PREGNANCY OR NURSING.
- **Do not shave the treated area the day of** or the day before the procedure to avoid irritation. Make sure the skin is shaved and hairless but do it at least 3 days before.

BLUE LIGHT FOR ACNE POST TREATMENT INSTRUCTIONS

- Until sensitivity has completely subsided, avoid all of the following:
 - Use of scented lotions or soaps, exfoliant creams (Retin-A, glycolic/salicylic and alphahydroxy acids), acne creams or gels, loofa sponges and aggressive scrubbing
 - o Hot or very cold water only wash with tepid lukewarm water
 - Shaving
 - Swimming pools and spas with chemicals/chlorine
 - Activities that cause excessive perspiration
- **Cold compresses**. A cold compress can be used to provide comfort if the treated area feels warm. This is rarely needed but if so use it within the first 24 hours after the treatment.
- Make Up. Makeup can be applied immediately after treatment as long as skin is intact and there are no signs of blistering.
- **Topical soothing products.** Vitamin E or Aloe Vera applied to the treatment area may provide a soothing effect.
- **Oral pain relievers**. Post treatment discomfort is typically minimal but if the area is very uncomfortable, oral pain relievers; i.e. Extra Strength Tylenol or Advil, may be used.
- Avoid sun exposure. Strictly avoid any sun exposure to the treated area for a minimum of 7 days after the procedure. The treated area is more prone to sunburn and pigmentation change. Keep the area covered and use an SPF of 30+, reapplying the sun block every two hours. Delayed blistering secondary to sun exposure has been noted up to 72 hours post treatment.
- Follow-up with recommended treatments. Subsequent treatments are based upon your clinician's recommendation and are typically 2 treatments per week for 2-6 weeks or until acne reduces. The number and frequency of treatments needed depend on the severity of your acne and how well it responds to treatment. Clinical studies suggest that at least two BBL 420 nm acne treatments are needed to begin to see any results and results have been reported to last up to 6 months after 4-6 treatments. Booster treatments are necessary as is a proper medical skincare regimen.

WHEN TO CONTACT US

If the skin is broken or blisters appears, apply an antibiotic ointment and contact the office immediately. Keep the affected area moist and avoid direct sunlight. Do not manipulate the area.

