

Buccal Fat Removal

Pre-Operative and Post-Operative Patient Instructions

PRE-OPERATIVE INSTRUCTIONS FOR BUCCAL FAT REMOVAL

SHOPPING LIST

☐ **Prescriptions**

Submitted to your pharmacy. Your pharmacy should contact you when ready to pick up.

☐ **Small Toothbrush**

Small toothbrush should be used for two weeks after surgery.

☐ **Tylenol (Acetaminophen):**

Alternative to narcotic pain medication if pain is not severe. Do NOT take NSAIDS.

☐ **Arnica**

Take 3 tablets 3 times a day starting 3 days before your procedure date.

NIGHT BEFORE SURGERY

- **DO NOT EAT OR DRINK ANYTHING AFTER MIDNIGHT OR SURGERY WILL BE CANCELLED.** If prescribed, take medications with a small sip of water.
- **Set Up Home Recovery Area:** This may include pillows, blankets, books, television, and anything else to assist with a comfortable recovery.
- **Bathing:** You can shower but do not apply lotion, perfume, hair product, etc.

DAY OF SURGERY

- **Dress Comfortably:** Dress in comfortable, clean, and loose-fitting clothes. Shirts that can be buttoned or zipped up are preferred.
- **DO NOT wear** any makeup, jewelry, cosmetic creams, hair products, deodorant, sunscreen, and remove all piercings.

POST-OPERATIVE INSTRUCTIONS FOR BUCCAL FAT REMOVAL

NIGHT OF SURGERY

- **Movement is Important:** Make sure you are up and walking around immediately after your surgery. When lying down in bed or on the couch, make sure you are moving your legs and ankles. Take deep breaths frequently to keep your lungs clear.



- **Sleep:** Sleep on your back with your head elevated about 30-40 degrees (2-3 pillows). Do not sleep on your side. Keeping your body more upright will minimize swelling. Continue this for one to two weeks.
- **Talking:** Avoid excessive talking, laughing, and chewing as this will cause more swelling and discomfort.

MEDICATIONS

- **Antibiotic:** Start your antibiotic tablet the night of surgery and take all tablets as prescribed and until finished.
- **Mouthwash:** Rise with any over the counter diluted mouthwash two to three times a day for one week. Dilute your mouthwash 1:1 with water.
- **Pain:** You will be prescribed a pain medication for post-operative pain control. If your discomfort after surgery is not strong you are welcomed to take Tylenol in place of the prescribed medication. Do not take Tylenol with the pain medication, often the medication you are prescribed will have Tylenol in it. **Do not exceed 4,000 mg of Tylenol in any 24-hour period.** Take medication with food to minimize risk of nausea.
- **Nausea:** A common side effect after general anesthesia and strong medications. If you are experiencing nausea, we advise that you take your nausea medication.
- **Constipation:** You will experience constipation if taking narcotic pain relievers. MiraLAX or other over the counter laxatives are recommended. Do not wait to take until you are constipated. Begin treatment with narcotic use.
- **Medications to Avoid:** Take only those medications approved or prescribed by your surgeon. Avoid medications containing aspirin or ibuprofen (Advil, Motrin, Naproxen, others) for two weeks before and after surgery. These medications may increase bleeding.
- **Substances to Avoid:** Avoid alcohol, nicotine, and caffeine, for these will dramatically slow the healing process.

NUTRITION

- **Diet:** Eat foods that are soft and require minimal chewing for one week. Do not eat salads, tough meats, or chew gum. Chewing can cause an increase in pain and swelling. Avoid sharp, granular foods or spicy foods.
- **Hydration:** Stay hydrated by drinking 8-10 glasses of water a day. Try to use straws when drinking fluids. Avoid alcohol while you are still taking pain medications.



BATHING

- **Showering:** You may wash your hair and face the next day after your surgery.
- **Hot Tubs/Baths/Swimming Pools:** No tub baths or Jacuzzi until your incisions have healed, and approved by your surgeon, which is usually around 2 weeks. It is best to wait one month for hot tubs as they tend to have more bacteria than regular chlorinated swimming pools.

HOW TO TAKE CARE OF YOUR INCISIONS

- **Eating:** AVOID opening your mouth more than 1 inch wide to prevent incision site separation (opening) for one week.
- **Cleaning your incisions:** rise your mouth with salt three times a day and after every meal. Continue this for one week.
- **Sutures:** All the sutures will be dissolvable.
- **Avoid brushing teeth for 48 hours.** For the next 2 weeks use a small toothbrush and be extra gentle.

WHAT TO EXPECT

- **Swelling and Discoloration:** It is completely normal to experience some swelling and discoloration from bruising around the surgical incision site.
- **Mild Bleeding:** Some mild bleeding is not unusual at the incision sites the first few days after surgery. If you try to do strenuous activity, increase your blood pressure by straining on the toilet, sneezing, heavy lifting or bending forward you will aggravate the bleeding and swelling.

ACTIVITY

- **Physical Activity:** It is important to get up, walk around, and not remain on complete bed rest while your body recovers. Normal daily activity may be resumed a few days after surgery. No bending over, straining, or lifting more than 5lbs for the first week after surgery.
- **Exercise:** Light physical activity may be resumed 2-3 weeks after surgery. Remember to start easy and build back up to your previous exercise levels. Just know that swelling may transiently be worse with exercise. Strenuous activity and contact sports should be avoided for the first week after surgery.

- **Driving:** Do NOT operate a vehicle or make important decisions until you have been off narcotic pain medications for 24 hours. Use good judgment.
- **Return to work:** Most patients return to work within 24-48 hours.
- **Cold Compress:** Use cool/moist compresses or ice wrapped in a small towel on the outside of your cheeks for 24-48 hours to reduce swelling. You may apply a compress for 20 minutes at a time and take a break every so often. **No Direct Ice!**

EMERGENCY SITUATIONS – WHEN TO CALL THE OFFICE (703-481-0002) OR GO TO THE HOSPITAL

- **Signs of Infection:** Spreading redness, worsening swelling, increased drainage or drainage of pus, worsening pain, warmth at incision site and temperature over 101 degrees Fahrenheit.
- **Excessive Bleeding:** If the incisions appear to be oozing blood excessively.
- **Other Emergency Situations:** Shortness of breath or difficulty breathing, chest pain, lightheadedness that does not quickly resolve, severe vomiting, pain, or asymmetric swelling in your legs