



Buccal Fat Pad Removal- Patient Instructions

Pre-Operative and Post-Operative Care | Outpatient Surgical Facility

PREPARATION BEFORE SURGERY

- Pick up and confirm you have all prescriptions before surgery.
- Purchase MiraLAX (stool softener), Tylenol (for pain), and cold compress supplies.
- Get Arnica tablets. If provided, take as directed starting after surgery.
- Nothing to eat or drink after midnight before surgery - if undergoing general anesthesia.
- Prepare a comfortable recovery area at home.
- Strictly follow and adhere to the “**Naderi Antibleeding Anti Bruising diet.**” See separate list.
- Avoid blood thinners and NSAIDs like aspirin, ibuprofen, naproxen, as well as nicotine, alcohol, and excessive caffeine.

DAY OF SURGERY

- Shower well. Shampoo hair and use antibacterial soap on face and body.
- Wear loose, cool, comfortable clothing. Button-up or zip-up shirts are preferred.
- Do not wear makeup, lotion, skin products, sun block, deodorant, or hair products
- Remove jewelry, piercings, contact lenses,
 - Remove hair extensions, weaves, braids and lash extensions (if possible) as they can harbor bacteria
- You will need an adult friend or relative to drive you home.

FIRST WEEK AFTER SURGERY

- Sleep on your back with your upper body elevated degrees. (Recliner or wedge pillow or multiple pillows)
- Apply cool compresses to your cheeks for 15–20 minutes every hour while awake for the first 3 days.
- Get up slowly and walk slowly but regularly with assistance to promote circulation.
- Avoid bending, straining, or lifting.
- You may feel weak and dehydrated so be careful of falling. Stay hydrated.
- You may shower 24 hours after surgery but be careful of falling.
 - Use gentle pressure and lukewarm water.

MEDICATIONS

- Take antibiotics as prescribed until finished. Notify us if you have any reactions.
- Use prescribed pain medication or Tylenol (if pain is mild).
- Do not exceed 4,000 mg of Tylenol in 24 hours.
- Begin stool softener when taking narcotic pain medication. Do not force or strain during bowel movement.
- Take your Medrol steroid dose pack as instructed.



INCISION CARE

- Use diluted Hydrogen Peroxide to rinse mouth gently 3x a day.
 - Mix half peroxide with half water in a small cup
- You may use a salt-water rinse instead.
 - Add 1 teaspoon kosher salt to one cup of room temperature water
- Do not pick or pull at incisions.
- Do not manipulate sutures with your tongue.
- All sutures are dissolvable.
- Use a small kid's toothbrush.
- **Avoid hard foods or seeds that may irritate or get stuck in the incision.**
- Avoid opening your mouth more than 1". Use a small spoon or small fork. Small bites.
- Soft foods and liquids are easier in the beginning.

ACTIVITY AND RECOVERY

- Avoid lifting over 20 pounds for 2 weeks.
- Most patients return to work in 1 week or sooner
- Avoid strenuous exercise for 3-4 weeks.
- Do not drive while taking narcotic pain medication.

NORMAL SYMPTOMS

- Swelling and bruising.
- Mild bleeding.
- Tightness, numbness, or mild itching around the area.
- Mild face or smile asymmetries.

NOTIFY YOUR DOCTOR IMMEDIATELY IF YOU EXPERIENCE:

- Fever greater than 101°F.
- Severe or worsening swelling.
- Excessive bleeding.
- Rash or severe itching or blisters.
- Call 911 if you experience shortness of breath or chest pain.
- Call 911 if having sudden increase in pain or swelling