



THE NADERI CENTER
PLASTIC SURGERY & DERMATOLOGY

Chin Implant

Chin Augmentation · Patient Instructions

Having more than one procedure? Follow the more cautious instruction. Direct instructions from Dr. Naderi always take precedence.

BEFORE YOUR SURGERY

- **Pick up your prescriptions well ahead of surgery.** We send them to your pharmacy at least 2 weeks in advance. Open the bag well ahead of your surgery and make sure nothing is missing. Call us right away if your antibiotics or pain medication are missing.

THE NADERI ANTI-BLEEDING, ANTI-BRUIISING DIET

Starting 2 weeks before surgery, follow this diet strictly. It directly affects how much you bleed during surgery and how much you bruise and swell afterward. Dr. Naderi considers it one of the most important parts of your preparation. See the separate Foods to Avoid list.

- Purchase MiraLAX (stool softener), Tylenol (for pain), hypochlorous acid (HOCl) spray, and cool compress supplies.
- Starting 1 week before surgery, spray hypochlorous acid (HOCl) on your chin and the skin beneath it 3 times a day as an antiseptic.
- Get Arnica tablets. Start them the day after surgery and take as directed.
- **Nothing to eat or drink after midnight the night before surgery,** or anesthesia will cancel your surgery.
- Prepare your recovery area at home with fluids, appropriate foods, and assistance.
- Starting 2 weeks before surgery, avoid aspirin, ibuprofen, naproxen, alcohol, and blood-thinning supplements. Check with your prescribing doctor before stopping any prescription blood thinner.
- Stop all nicotine (smoking, vaping, patches, gum) at least 2 weeks before surgery; 4 weeks is preferred.
- Moderate caffeine is fine; one cup of coffee or tea is acceptable.
- Make sure your dental hygiene is good. Problems with your teeth can lead to an infected chin implant.

- For men: shave the day before surgery. No facial hair and no beard.

THE NIGHT BEFORE & DAY OF SURGERY

- **Take two showers: one the night before surgery and one the morning of surgery.** Each time, wash your hair well and use antibacterial soap on your face and body. Men, make sure you are shaved.
- Wear loose, comfortable clothing that buttons or zips in the front.
- Arrive without makeup, lotions, skin products, sunblock, deodorant, or hair products.
- Remove jewelry and piercings. Do not wear contact lenses; wear your glasses instead.
- Remove hair extensions, weaves, braids, and lash extensions if possible. They can harbor bacteria.
- Arrange for a responsible adult to drive you home and stay with you initially. **You will not be released to an Uber or taxi.**

THE FIRST WEEK

- Sleep on your back with your upper body elevated about 45° (recliner, wedge pillow, or several pillows).
- Use cool compresses or frozen peas wrapped in cloth for 15–20 minutes about every hour while awake for the first 3 days.
- **Never place ice or frozen materials directly on the skin.** Numb skin can get frostbite.
- Rest, hydrate, and walk gently and regularly to promote circulation. Rise slowly and have assistance at first; you may feel weak.
- Avoid bending, straining, or lifting.
- Eat fresh pineapple, including the center core, or drink cold-pressed pineapple juice. Its bromelain helps reduce swelling.
- **Take your antibiotics.** They are extremely important in preventing infection.

- **You may shower 48 hours after surgery.** Use lukewarm water and gentle pressure. Do not let the spray hit your incisions directly; let the water run over them gently. Do not rub your incisions; fresh incisions can open if handled roughly. Be careful of falling.
- **Do not press on or move the implant.**

MEDICATIONS

- Start your antibiotics the evening of surgery, once your nausea has passed and you can keep food down. Take them as prescribed until finished, and tell us if you have any reaction.
- Use your prescribed pain medication, or Tylenol if pain is mild. Do not take the two together; most prescription pain medications already contain Tylenol.
- Do not exceed 4,000 mg of Tylenol (acetaminophen) in 24 hours.
- Use your nausea medication as needed, if prescribed.
- Begin a stool softener when you start narcotic pain medication. Do not strain during bowel movements.
- Start your Medrol steroid dose pack the day after surgery: 6 pills on the first day, then 5, 4, 3, 2, and 1 on the days that follow, until finished. If your pharmacy gives you the pills loose in a bottle, follow the same schedule.

INCISION CARE

- Once Dr. Naderi directs you on your telehealth visit, remove your pressure wrap the afternoon after surgery.
- You will most likely have tape covering the incision. Leave it alone. If it falls off, clean the incision as described below.
- Clean incisions gently 3–4 times a day with a Q-tip dipped in diluted hydrogen peroxide. Roll the Q-tip; do not scrub.
 - Mix half peroxide and half water in the cap, then discard the excess.
- Apply prescribed antibiotic ointment or Vaseline very gently 3–4 times a day.
- Continue your hypochlorous acid (HOCl) spray on your incision and the surrounding skin 3 times a day.
- Brush your teeth with a small toothbrush and use a mouth rinse 2–3 times a day to keep your mouth clean.
 - Be gentle.
 - No flossing.
- Do not pick or pull at incisions.
- Dr. Naderi typically removes sutures 1 week after surgery.
- Protect healing skin and incisions from direct sun exposure for 3 months.

ACTIVITY AND RECOVERY

- Do not lift more than 20 pounds for the first month.
- No exercise for the first month. You may then gradually resume and build back slowly, with the goal of returning to full activity by about 3 months.
- Most patients return to work in 1 week.
- Do not drive while taking narcotic pain medication or while you feel impaired.
- Do not fly for 7–10 days after surgery. Flying can increase swelling for the first few months, so avoid unnecessary flights during that time.

WHAT IS NORMAL

- Swelling and bruising
- Stiffness, tightness, and a restricted smile
- Mild asymmetries of the chin and smile
- A noticeable incision early on

If you are ever unsure, Dr. Naderi and our team are always available. Take a few clear, well-lit photos and email them to Surgery@NaderiCenter.com. For anything more urgent, please call the office so we get your message quickly.

CALL US RIGHT AWAY IF YOU HAVE

- Fever over 101°F
- Rapidly increasing or severe swelling
- Worsening pain not improving with medication
- Bleeding, increasing redness, warmth, pus, or drainage
- Rash, severe itching, or blisters (possible medication reaction)

CALL 911 IMMEDIATELY FOR

- Chest pain
- Shortness of breath
- Excessive bleeding

REACH US

VIRGINIA 703.481.0002

MARYLAND 301.222.2020

EMAIL Surgery@NaderiCenter.com

WEB www.NaderiCenter.com